

MATHEMATICS ENRICHMENT CLUB.
Solution Sheet 8, July 2, 2018

1. The digital sum of 13950264876 is 51. Since $51 = 17 \times 3$, it is a multiple of three, but it is not a multiple of nine. Thus the prime factorisation of 13950264876 contains 3, but not 3^2 . Thus it cannot be a square.
2. Let the integer we are looking for be n and the common remainder be r . Then there are integers a , b and c such that $364 = an + r$, $414 = bn + r$ and $539 = cn + r$. But then $414 - 364 = (b - a)n$, $539 - 414 = (c - b)n$ and $539 - 364 = (c - a)n$, so we can see that n divides the differences of the three numbers. Now


$$414 - 364 = 50$$

$$539 - 414 = 125$$

$$539 - 364 = 175$$

The largest number that divides all three differences is 25.

3. We will first show that $\triangle AEF$ is similar to $\triangle ACB$, with a scale factor of $\frac{1}{2}$.



Since E is the midpoint of AC , $AE = \frac{1}{2}AC$. Similarly, $AF = \frac{1}{2}AB$. Furthermore, $\angle A$ is common to both triangles. So $\triangle AEF \sim \triangle ACB$, (ASS), with a scale factor of $\frac{1}{2}$. As a result, $EF = \frac{1}{2}BC$, and $\angle AEF = \angle ACB$. Thus $EF \parallel BC$.

4. Since acute angled triangles exist, we know that there are convex polygons with at least three acute angles.

If the interior angle of a polygon is acute, then the exterior angle must be obtuse. The sum of the exterior angles of a convex polygon is 360° . As the sum of four numbers greater than 90° is greater than 360° , a convex polygon must have less than four acute angles. Thus three is the largest number of acute angles that a convex polygon can have.

5. Let a and b be integers such that $a^3 = x + \sqrt{x^2 + 1}$ and $b^3 = x - \sqrt{x^2 + 1}$. Then $\frac{a^3 + b^3}{x + \sqrt{x^2 + 1} + x - \sqrt{x^2 + 1}} = a + b = y$, where $y \in \mathbb{Z}$. Now

$$(a + b)^3 = a^3 + b^3 + 3ab(a + b);$$

but

$$\begin{aligned} a^3 + b^3 &= x + \sqrt{x^2 + 1} + x - \sqrt{x^2 + 1} = 2x \\ ab &= \frac{(x + \sqrt{x^2 + 1})(x - \sqrt{x^2 + 1})}{x^2 - (x^2 + 1)} \\ &= \frac{-1}{x^2 - (x^2 + 1)} \\ &= \frac{-1}{-1} = 1; \end{aligned}$$

Consequently,

$$\begin{aligned} y^3 &= 2x + 3y \\ \Rightarrow x &= \frac{y^3 + 3y}{2}; \quad y \in \mathbb{Z} \end{aligned}$$

We can see that y^3 and $3y$ have the same parity, and so x is an integer.

6. (a) $\phi(12) = 4$ and $\phi(30) = 8$.
- (b) We can think of $\phi(n)$ as being the number of positive integers less than n which are not a multiple of a factor of n (except the factor 1). So if p is prime, its only factors are 1 and p . Thus $\phi(p) = p - 1$.
For p^2 , the factors are 1, p and p^2 , so the multiples of the factors that aren't 1 are $p; 2p; 3p; \dots; p^2$, of which there are p . So $\phi(p^2) = p^2 - p = p(p - 1)$.
For p^3 , the factors are 1, p , p^2 and p^3 . Multiples of the factors that aren't 1 are $p; 2p; 3p; \dots; p^2, (p + 1)p; \dots; 2p^2; \dots; p^3$. That is, a total of p^2 factors. So $\phi(p^3) = p^3 - p^2 = p^2(p - 1)$.
- (c) Using the same method as above, the factors of pq are 1, p , q and pq . The multiples of the factors that aren't 1 are $p; 2p; \dots; qp$ (q multiples) and $q; 2q; \dots; pq$ (p multiples), but we don't want to count pq twice. So $\phi(pq) = pq - q - p + 1 = (p - 1)(q - 1)$.

Senior Questions

1. (a) Expanding the right hand side,

$$\begin{aligned}(n^2 - 3n - 1)^2 - 25n^2 &= n^4 - 2n^2(3n + 1) + (3n + 1)^2 - 25n^2 \\ &= n^4 - 6n^3 - 2n^2 + 9n^2 + 6n + 1 - 25n^2 \\ &= n^4 - 6n^3 - 18n^2 + 6n + 1\end{aligned}$$

(b) We can use the previous result to factorise $n^4 - 6n^3 - 18n^2 + 6n + 1$. So

$$(n^2 - 3n - 1)^2 - 25n^2 = [(n^2 - 3n - 1) - 5n][(n^2 - 3n - 1) + 5n]$$

Using the symmetry of the graph, we can estimate that the largest root is

$$x_{max} = \frac{11}{2} + \frac{1}{2} x_5 \quad 17:763552537181550$$

$$f(x_{max}) = 3.55 \cdot 10^{-15}$$

3. Let ABC be a triangle, and let D , E and H be the midpoints of BC , AC and AB , respectively. Suppose that O is the point of intersection of BE and AD . Let F and G be the midpoints of OA and OB , respectively. Then, applying the mid-line theorem to $\triangle AOB$, $FG \parallel AB$, and $FG = \frac{1}{2}AB$. Similarly, by applying the mid-line theorem to $\triangle ACB$, we can see that $ED = \frac{1}{2}AB$ and $ED \parallel AB$. Thus $DEFG$ is a parallelogram, and O is the point of intersection of its two diagonals. Thus $OD = OF = AF$ and $OE = OG = GB$. Consequently, O is located $\frac{1}{3}$ the way along the medians AD and BE from their respective feet.

By a similar argument, we can show that the point of intersection of the medians CH and AD lies $\frac{1}{3}$ the length of AD away from D . Thus the two points of intersection coincide, and the three medians are concurrent.

