

MV²⁺. Likewise, Tris-HCl buffer (25 mM, pH 7.5 with 100 mM KCl) solutions of GroEL-CdS complex and cpn-CdS complex (2 ml, 700 nm based on Cd²⁺) were titrated. Emission ($\lambda_{\text{ext}} = 370 \text{ nm}$) spectra were recorded on a FP-777W spectrophotometer (JASCO).

Thermal stability

Fluorescence spectra ($\lambda_{\text{ext}} = 370 \text{ nm}$, wavelength of observed fluorescence $\lambda_{\text{obsd}} = 530 \text{ nm}$) of GroEL-CdS complexes and cpn-CdS complexes were recorded at designated temperatures on a FP-777W spectrophotometer (JASCO), where the fluorescence intensities at 4 °C were used as the bases for relative fluorescence intensities. The temperature was directly controlled by a ECT271 Peltier thermometric apparatus (JASCO; 40 °C min⁻¹ on heating and 25 °C min⁻¹ on cooling).

ATP response

To a 2-ml Tris-HCl buffer (25 mM, pH 7.5 with 100 mM KCl) solution of cpn-CdS complexes (0.5 μM based on cpn) were added aqueous solutions of ATP (100 mM) and MgCl₂ (1 M) ([ATP] = 20 μM , [Mg²⁺] = 25 mM after mixing), and the mixture was incubated at 70 °C for 10 min. The supernatant solution was subjected to fluorescence spectroscopy and analytical SEC with an UV/fluorescence dual detector.

Received 11 November 2002; accepted 14 April 2003; doi:10.1038/nature01663.

- Alivisatos, A. P. Semiconductor clusters, nanocrystals, and quantum dots. *Science* **271**, 933–937 (1996).
- Schmid, G. et al. Current and future applications of nanoclusters. *Chem Mater* **28**, 179–185 (1999).
- Meldrum, F. C., Heywood, B. R. & Mann, S. Magnetoferritin — in vitro synthesis of a novel magnetic protein. *Science* **257**, 522–523 (1992).
- Wong, K. K. W. & Mann, S. Biomimetic synthesis of cadmium sulfide-ferritin nanocomposites. *Angew Chem Int Ed Engl* **8**, 928–932 (1996).
- Shenton, W., Pum, D., Sleytr, U. B. & Mann, S. Biocrystal templating of CdS superlattices using self-assembled bacterial S-layers. *Science* **277**, 585–587 (1997).
- Balogh, L. & Tomalia, D. A. Poly(amidoamine) dendrimer-templated nanocomposites. I. Synthesis of zerovalent copper nanoclusters. *Chem Mater* **10**, 7355–7356 (1998).
- Lemon, B. I. & Crooks, R. M. Preparation and characterization of dendrimer-encapsulated CdS semiconductor quantum dots. *Chem Mater* **12**, 12886–12887 (2000).
- Shenton, W., Mann, S., Colfen, H., Bacher, A. & Fischer, M. Synthesis of nanophase iron oxide in lumazine synthase capsids. *Science* **291**, 442–445 (2001).
- McMillan, R. A. et al. Ordered nanoparticle arrays formed on engineered chaperonin protein templates. *Science* **297**, 247–252 (2002).
- Roseman, A. M., Chen, S., White, H., Braig, K. & Saibil, H. R. The chaperonin ATPase cycle: mechanism of allosteric switching and movements of substrate-binding domains in groEL. *Cell* **87**, 241–251 (1996).
- Ranson, N. A. et al. ATP-bound states of groEL captured by cryo-electron microscopy. *Cell* **107**, 869–879 (2001).
- Hendrix, R. W. Purification and properties of groE, a host protein involved in bacteriophage assembly. *J Mol Biol* **129**, 375–392 (1979).
- Braig, K. et al. The crystal structure of the bacterial chaperonin groEL at 2.8 Å. *Science* **267**, 578–586 (1994).
- Taguchi, H., Konishi, J., Ishii, N. & Yoshida, M. A chaperonin from a thermophilic bacterium, *Thermoplasma acidophilum*, that controls refoldings of several thermophilic enzymes. *J Biol Chem* **266**, 22411–22418 (1991).
- Amada, K. et al. Molecular cloning, expression, and characterization of chaperonin-60 and chaperonin-10 from a thermophilic bacterium, *Thermoplasma acidophilum* HB8. *J Biol Chem* **270**, 347–354 (1995).
- Ishii, N., Taguchi, H., Sumi, M. & Yoshida, M. Structure of holo-chaperonin studied with electron microscopy. Oligomeric cpn10 on top of two layers of cpn60 rings with two stripes each. *FEBS Lett* **299**, 169–174 (1992).
- Ishii, N., Taguchi, H., Sasabe, H. & Yoshida, M. Folding intermediate binds to the bottom of bullet-shaped holo-chaperonin and is readily accessible to antibody. *J Mol Biol* **236**, 691–696 (1994).
- Murakoshi, K. et al. Preparation of size-controlled hexagonal CdS nanocrystallites and the characteristics of their surface structures. *Chem Mater* **10**, 579–586 (1998).
- Hosokawa, H. et al. EXAFS observation of the surface structure of colloidal CdS nanocrystallites in N,N-dimethylformamide. *J Phys Chem* **100**, 6649–6656 (1996).
- Brus, L. E. Electron-electron and electron-hole interactions in small semiconductor crystallites: The size dependence of the lowest excited electronic state. *J Chem Phys* **80**, 4403–4409 (1984).
- Ramsden, J. J. & Grätzel, M. Photoluminescence of small cadmium sulfide particles. *J Chem Phys* **80**, 919–933 (1984).
- Henglein, A. Photochemistry of colloidal cadmium sulfide. 2. Effects of adsorbed methyl viologen and of colloidal platinum. *J Phys Chem* **86**, 2291–2293 (1982).
- Llorca, O., Galán, A., Carrascosa, J. L., Muga, A. & Valpuesta, J. M. GroEL under heat-shock. *J Biol Chem* **273**, 32587–32594 (1998).
- Martin, J., Horwich, A. L. & Hartl, F. U. Prevention of protein denaturation under heat stress by the chaperonin Hsp60. *Science* **258**, 995–998 (1992).
- Goloubinoff, P., Christeller, J. T., Gatenby, A. A. & Lorimer, G. H. Reconstitution of active dimeric ribulose biphosphate carboxylase from an unfolded state depends on two chaperonin proteins and Mg-ATP. *Science* **242**, 884–889 (1989).
- Todd, M. J., Viitanen, P. V. & Lorimer, G. H. Hydrolysis of adenosine 5'-triphosphate by *Escherichia coli*

($Fe_{HR} = Fe_D + Fe_P$) and Fe_T , and the ratio $Fe_P/(Fe_P + Fe_H)$, known as the degree of pyritization (DOP), have been used successfully to evaluate the redox state of ancient oceans¹⁰⁻¹³. DOP < 0.45 and $Fe_{HR}/Fe_T < 0.38$ are generally found for sediments depositing from oxic bottom water, whereas DOP > 0.45 and $Fe_{HR}/Fe_T > 0.38$ may be diagnostic for euxinic sediments^{10,12}. The enrichment of Fe_{HR} and high DOP in euxinic sediments arises fundamentally from

sea sediments are comparable in magnitude, but oxic environments record a much greater range of values^{13,20}.

The same is true for Phanerozoic examples. Gauthier²¹ measured $\delta^{34}\text{S}$ values for pyrites deposited beneath oxic and anoxic bottom waters of the Cretaceous seas that flooded North America. Like our Roper samples,

The documentation of euxinic and low sulphate conditions in mid-Proterozoic marine basins paves the way to an improved understanding of early life and environments. In such oceans, methanogenic archaeans could have played an enhanced role in the carbon cycle, contributing to long-lived greenhouse conditions²⁷. Low sulphate may also help to explain the prominence of penecontemporaneous dolomite in mid-Proterozoic and older carbonate platforms²⁸. Through its effects on biologically important trace elements, seawater chemistry may help to explain the ecological and evolutionary distributions of early eukaryotic photoautotrophs⁸. And, if sulphate levels remained low until the latest Proterozoic, oxygen probably also remained well below present levels, influencing the early diversification of animals²⁹. Further biogeochemical research carried out in the framework of sequence stratigraphy